

NONTHERMAL RADIO EMISSION FROM PLANETARY NEBULAE

Ruth Dgani

knill@astro.as.utexas.edu

Department of Astronomy, The University of Texas

Austin, TX 78712

and

Noam Soker

soker@physics.technion.ac.il

Department of Physics, University of Haifa at Oranim

Tivon 36006, Israel

ABSTRACT

In a recent analysis of the radio emission from the planetary nebula A30, Dgani, Evans & White (1998) claim that the emission, located in the inner region, is probably dominated by nonthermal emission. We propose a model to explain this. We assume that the fast wind, blown by the central star of A30 carries a very weak magnetic field. The interaction of this wind with a cluster of dense condensations traps the magnetic field lines for a long time and stretches them, leading to a strong magnetic field. If relativistic particles are formed as the fast wind is shocked, then the enhanced magnetic field will result in nonthermal radio emission. The typical nonthermal radio flux at 1 GHz can be up to several milli-Jansky. In order to detect the nonthermal emission, the emitting region should be spatially resolved from the main optical nebula. We list other planetary nebulae which may possess nonthermal radio emission.

Subject heading: ISM: magnetic fields – ISM: kinematics and dynamics – Radio continuum: ISM – Planetary nebulae: general – Planetary nebulae: individual (A30)

1. INTRODUCTION

The possibility of magnetic fields carried by the fast winds in planetary nebulae (PNs; Chevalier and Luo 1994; hereafter CL; Chevalier 1995; Garcia-Segura 1997) raises the question of whether the interaction between this wind and the previously ejected slow wind gives rise to particle acceleration, and hence nonthermal emission from relativistic electrons (Chevalier 1995). Chevalier notes that the major problem with detecting the nonthermal emission is the strong thermal free-free emission emanating from the dense shell. In many PNs, however, knots and tails are present relatively close to the central star, where both the ram pressure of the fast wind and the magnetic field are higher, increasing the nonthermal emission. In the hydrogen deficient PNs A30 and A78, for example, knots with tails appear in a disk structure close to the central star (Borkowski, Harrington, & Tsvetanov 1995). These nebulae are believed to have had a helium flash late in their evolution, and are known to possess fast winds, $v_w \sim 4000 \text{ km s}^{-1}$ (Leuenhagen, Koesterke & Hamann 1993). The knots are believed to be photo evaporating globules (Dyson 1968; Kahn 1969; Borkowski *et al.* 1995). The HST images of the knots in the central region of A30 (Borkowski *et al.* 1995) show that they are being disintegrated and accelerated by the dilute shocked fast wind from the central star, giving rise to Rayleigh-Taylor instability. In addition, hydrogen deficient nebulae have lower thermal emission (see discussion in Dgani, Evans & White 1998), allowing the nonthermal component to be detected.

Jones, Kang & Tregillis (1994) have simulated the evolution of dense gas clouds moving supersonically through a magnetized low density medium. They took a passive magnetic field and found that the synchrotron emission from relativistic electrons increase rapidly when the cloud begins to fragment under Rayleigh-Taylor instability. The synchrotron emission is dominated by field enhancement through stretching in the shear turbulent layers formed between and behind the fragments. Jones, Ryu & Tregillis (1996) performed full two dimensional MHD calculations and found that when the ambient field is transverse to the flow fast amplification of the field occurs also at the nose of the cloud, and not only in the turbulent regions.

Dgani *et al.* (1998) present a radio map of the central region of A30. The map shows an

extended source of about 0.15 mJy stretching from the central star to a point 3" to the west. Based on a comparison to the optical line emission from that region, they claim that the source is probably nonthermal. Motivated by this recent finding, we apply the theoretical results of Jones *et al.* (1994; 1996) to the interaction of a magnetic fast wind with knots in PNs. The amplification of the magnetic field is calculated in §2, while in §3 we estimate the nonthermal radio emission from the interaction region. In §4 we summarize, and list PNs, other than A30 (Dgani *et al.* 1998), which we expect to have nonthermal radio emission.

2. INTERACTION OF DENSE KNOTS WITH A MAGNETIZED WIND

Like CL, we examine a slowly rotating central star, for which the tangential component of the magnetic field dominates far from the star. For a field line that starts radially from the stellar equator, the downstream tangential component of the field at $r \gg r_s$, where r is the distance from the central star and r_s its radius, is given by

$$B = B_s \frac{v_{\text{rot}}}{v_w} \frac{r_s}{r}, \quad (1)$$

where v_{rot} is the equatorial velocity of the rotating star, v_w is the fast wind terminal velocity, and B_s is the (radial) magnetic field on the stellar surface. Since both the magnetic pressure $B^2/8\pi$ and the ram pressure ρv_w^2 drop as r^{-2} before hitting the shock wave, it is useful to define the parameter (CL):

$$\sigma = \frac{B^2}{4\pi\rho v_w^2} = \frac{B_s^2 r_s^2}{\dot{M} v_w} \left(\frac{v_{\text{rot}}}{v_w} \right)^2, \quad (2)$$

where $\dot{M}$ is the mass loss rate to the fast wind and $\rho = \dot{M}(4\pi r^2 v_w)^{-1}$ is the density in the wind. σ is the ratio of field energy density to kinetic energy density in the wind. CL find that the magnetic field shapes the nebula for $\sigma \gtrsim 10^{-4}$. From equation (2) we can write the magnetic field in the equatorial plane in the free wind as

$$B = 3.5 \times 10^{-7} \left(\frac{\sigma}{10^{-8}} \right)^{\frac{1}{2}} \left(\frac{\dot{M}}{5 \times 10^{-8} M_\odot \text{ yr}^{-1}} \right)^{\frac{1}{2}} \left(\frac{v_w}{4000 \text{ km s}^{-1}} \right)^{\frac{1}{2}} \left(\frac{r}{10^{16} \text{ cm}} \right)^{-1} G. \quad (3)$$

The mass loss rate is scaled according to the value Borkowski *et al.* (1995) used for A30.

Consider first a single globule with radius R_g . A supersonic flow incident on it will form a bow shock at its nose, a turbulent shear layer around its sides and a turbulent wake behind it. The magnetic field is expected to be enhanced in these regions because of line stretching (e.g., Jones *et al.* 1994, 1996). As shown in the extensive numerical simulations of Jones *et al.* (1996), when the magnetic field lines are transverse to the flow the magnetic pressure on the nose of the clump reaches equipartition with the ram pressure, even if the initial magnetic field is quite weak. Using their analysis for the numerical results, we can estimate the growth rate of the magnetic field by the following consideration (their equations 8-10)

Jones *et al.* (1996, eq. 8) start with the ideal MHD magnetic induction equation

$$\frac{d\ln(B/\rho)}{dt} = \frac{B \cdot [(B \cdot \nabla)u]}{B^2}. \quad (4)$$

u , ρ and B are the velocity, the density and the magnetic field in the flow respectively. Taking $(B \cdot \nabla)u \sim Bu/R_g$, equation (4) becomes

$$\frac{d\ln(B/\rho)}{dt} = \frac{u}{R_g}. \quad (5)$$

The last equation gives exponential growth to the magnetic field as long as the magnetic field lines stay near the stagnation point. Jones *et al.* (1996) estimate that at the stagnation point the magnetic pressure will rise exponentially until the bullet is crushed or until the magnetic pressure equals the ram pressure. However, since they performed 2D calculations, they neglected the sliding of the magnetic field line around the bullet. Although they claim irregularities in the 3D spherical bullet will capture the field lines and prevent sliding, we believe that for a flow with very large $\beta = P_{\text{th}}/P_B$, the field lines will be carried around the globule in a time $\lesssim R_g/u$, much shorter than the bullet's crushing time. Here P_{th} and P_B are thermal and magnetic pressures, respectively.

Other sites of magnetic field enhancement are the turbulent mixing layer at the sides of the globule, and its turbulent wake. These are more promising sites, mainly because the clumps (globules) appear in groups, and turbulent flows dominate the region.

In here, we consider the flow around and inside an ensemble of clumps, occupying a region of size R_{ens} . The typical distance between the clumps is of the order of their size R_g . This case has not been studied numerically, but we can apply an analytical approach to estimate the magnetic field enhancement. A turbulent flow will form between the globules, where the magnetic field can be amplified. There, the change of the wind velocity u along a field line depends on the size of the mixing layer $l \sim R_g$. That is $(B \cdot \nabla)u \sim Bu/R_g$. The time a field line spends in the turbulent region is the time it takes the flow to pass the group: R_{ens}/u . Substituting these values in the induction equation (4), and integrating over the time the field lines spend inside the group of globules, gives the relative amplification

$$B/\rho \sim \exp \frac{R_{\text{ens}}}{R_g}. \quad (6)$$

For $R_g \sim 0.1R_{\text{ens}}$, for example, there is a large amplification factor $\sim 10^4$. Therefore, the magnetic field will reach equipartition even if its initial value is $\sim 10^{-4}$ that of equipartition, i.e., $\sigma \sim 10^{-8}$ (eq. 2). In the model of CL and Garcia-Segura (1997) the post shock magnetic field grows as $B \sim r$, which gives a factor of $\sim 10 - 100$ for the magnetic field and $\sim 100 - 10^4$ for the magnetic pressure. Since CL assume that the magnetic field shapes the PN globally, they have to assume that the initial magnetic pressure is $\gtrsim 10^{-4}\rho v^2$, where ρv^2 is the fast wind's ram pressure. This means $\sigma > 10^{-4}$. In the flow pattern assumed here, the localized turbulence results in a much larger increase in the magnetic field. Therefore, to achieve equipartition between the magnetic pressure and the post shock thermal pressure, a value of $\sigma \gtrsim 10^{-8}$ is sufficient.

3. NONTHERMAL RADIO EMISSION

The upper limit for the enhanced magnetic field is the equipartition value, found from the relation $P_B = \rho v_w^2$,

$$B_{\text{max}} \simeq 5 \times 10^{-3} \left(\frac{\dot{M}}{5 \times 10^{-8} M_{\odot} \text{ yr}^{-1}} \right)^{\frac{1}{2}} \left(\frac{v_w}{4000 \text{ km s}^{-1}} \right)^{\frac{1}{2}} \left(\frac{d}{10^{16} \text{ cm}} \right)^{-1} G, \quad (7)$$

where d is the distance between the central star and the group of globules. If the relativistic electrons are produced in the bow shock and then carried downstream to the magnetized region,

they can emit synchrotron radiation. We assume equipartition between the magnetic field and relativistic particles (electrons and nuclei). The radio luminosity between ν_1 and ν_2 can then be estimated if the spectral index α , (i.e., the flux $F_\nu \propto \nu^{-\alpha}$), of the radiation is known (cf. Pacholczyk 1970 Page 171, Eq. 7.14). The radio luminosity is given by

$$L = \frac{B^{7/2} \Phi R_{ens}^3}{4.5 (1 + k) c_{12}(\alpha, \nu_1, \nu_2)}, \quad (8)$$

where Φ is the fraction of the ensemble volume occupied by magnetic field and relativistic particles, k is the ratio of the heavy particle energy to the electron energy, and $c_{12}(\alpha, \nu_1, \nu_2)$ is tabulated in appendix 2 of Pacholczyk.

Taking $\nu_1 = 10^7$ Hz and $\nu_2 = 10^{11}$ Hz, $\alpha = 0.75$, for which $c_{12} = 5 \times 10^7$ and $k = 100$, (Chevalier 1977;1995) we obtain

$$L \simeq 4.9 \times 10^{28} \Phi \left(\frac{R_{ens}}{5 \times 10^{15} \text{cm}} \right)^3 \left(\frac{B}{5 \times 10^{-3} G} \right)^{7/2} \text{erg s}^{-1}. \quad (9)$$

The radio flux at 1GHz can now be estimated:

$$F_{\nu 9} \simeq 3.5 \Phi \left(\frac{R_{ens}}{5 \times 10^{15} \text{cm}} \right)^3 \left(\frac{B}{5 \times 10^{-3} G} \right)^{7/2} \left(\frac{D}{kpc} \right)^{-2} \text{mJy} \quad (10)$$

where D is the distance to the source, and $1 \text{mJy} = 10^{-29} \text{W m}^{-2} \text{Hz}^{-1}$. A few mJy is much less than the observed fluxes from the shells of most catalogued PNs (Zijlstra, Pottasch, & Bignell 1989). However, the emission we predict is far from the PN shell, near the central star, and will not be overwhelmed by thermal emission. Chevalier (1995) discussed the case in which nonthermal electrons are produced in the fast wind termination shock, and then carried downstream to the magnetized region near the shell. In this case the nonthermal emission region neighbors the thermal emission region and cannot be separated observationally. Therefore, Chevalier concluded that a search for radio synchrotron emission from planetary nebulae would involve a frequency lower than 1GHz.

Dgani *et al.* (1998) show an 8 GHz radio map of the hydrogen deficient PN A30. They present a source of ~ 0.1 mJy which extends to about $3''$ west of the central star. A30 possesses a large shell ($120''$) visible in both H_α and OIII, and a hydrogen deficient central region visible

in OIII but not in H_α (Jacoby 1979). The radio emission emanates from the central region. The source extends to the vicinity of the main blobs of the HST image (Borkowski *et al.* 1995), but it cannot be identified with any of them. This is compatible with our model where the radio emission results from magnetic amplification in the turbulent region formed by decelerating clumps.

A somewhat parallel situation occurs in young SNRs. In the SNR Cassiopea A (Cas A), the radio knots as well as the optical knots appear to be connected to the clumpy component of the ejecta, however, they are not identical. Anderson *et al.* (1994) analyze the observations of optical and radio emission knots in the SNR Cas A. They claim that while the clumps are almost certainly sites of particle acceleration, when there is substantial magnetic field growth, it will enhance the synchrotron emission. Complex flow patterns bring in particles that were accelerated in different regions. In young SNRs particles are produced in many acceleration sites: the blast wave shock, the reverse shock and many internal interactions shocks. Thus the enhanced emission in the radio knots is probably a result of field amplification. The numerical simulations of Jun & Jones (1998) and of Jun, Jones & Norman (1996) support this conclusion.

4. DISCUSSION AND SUMMARY

We studied the interaction of a weakly magnetized fast wind, of velocity $v_w > 1,000 \text{ kms}^{-1}$, blown by the central stars of PNs with high density condensations, and we propose that it leads to the amplification of the magnetic field. When this interaction is with a cluster of clumps, the magnetic field lines are trapped for a long time in the interaction region. This process is equivalent to the numerical simulations of Jones *et al.* (1996), in which the 2D nature of the flow did not allow the magnetic field lines to slide around the condensation. Jones *et al.* (1996) found that the magnetic field can reach equipartition with the wind’s ram pressure.

The enhanced magnetic field will have several effects: (1) It will reduce the mixing (entrainment) of the clump material with the fast wind. (2) It will reduce the heat conduction between the cool clump and the hot shocked fast wind material. (3) The magnetic field lines will

reconnect among themselves. This process may lead to heating and further X-ray emission. (4) It may result in nonthermal radio emission. The first three processes will affect the X-ray emission from the hot material. In the current study we did not examine these processes. We explored here, and concluded that the expected nonthermal radio emission can be detected in PNs with fast winds and clumps close to the central star. Such a PN is A30 (Dgani *et al.* 1998), claimed also to have extended X-ray emission partially overlap with the location of the optical condensation (Chu, Chang & Conway 1997).

The required initial magnetic field for detectable nonthermal radio emission is ~ 2 orders of magnitude lower than the field intensity required to influence the shape of PNs, according to the model of CL and Garcia-Segura (1997). The stronger magnetic field required by CL and Garcia-Segura (1997) requires much faster rotation, as shown by equation (2) with $\sigma > 10^{-4}$. This is very problematic for their model, since the central stars of PNs are expected to rotate slowly because of mass loss. This is true even if their AGB progenitors are spun-up by companions on the AGB (Soker 1998). The field in the model of CL and Garcia-Segura (1997) is globally large. The amplification of the magnetic field through the interaction of the magnetized wind with a cluster of condensations, on the other hand, enhances the field locally, but not globally. Therefore, the detection of strong magnetic fields in specific locations in PNs does not necessarily mean that the magnetic field is globally strong.

The process proposed in this study predicts detectable radio synchrotron emission from PNs having fast winds and high mass loss rates, both of which determine the magnetic field intensity (eq. 7), and optical condensations. In order to detect this synchrotron emission we need to resolve the central region of the nebula from the shell. This requires high spatial resolution observations.

To list several PNs, in addition to A30 and the other hydrogen deficient PNs (see discussion in Dgani *et al.* 1998), for which we expect nonthermal radio emission we select PNs having fast winds from Patriarchi & Perinotto (1991) and check their optical images for condensations in the inner region. We find that from the list of Patriarchi & Perinotto (1991) only NGC 2371 and NGC 7094 show no structure in the very inner regions. Therefore, in general all PNs with fast winds

blown by their central stars should be searched for nonthermal emission, in particular, those with strong condensations in their central regions: A78, NGC 5189 and NGC 6210.

ACKNOWLEDGMENTS: It is a pleasure to thank Kazik Borkowski, Patrick Harrington, Neal Evans, Mordecai Mac-Low, and Stephen White, for useful discussions. We thank the referee Tom Jones for insightful suggestions. We thank Sarah Baird, Neal Evans and Stephen White for a careful reading of the manuscript and useful comments. This research has been supported in part by a grant from the Israel Science Foundation.

REFERENCES

- Anderson, M.C., Jones, T.W., Rudnick, L., Tregillis, I.L., & Kang, H. 1994, ApJ, 421, L31.
- Borkowski, K.J., Harrington, J.P., & Tsvetanov, Z.I., 1995, ApJ, 449, L143.
- Chevalier, R. A. 1977, A&A, 59, 289
- Chevalier, R. A. 1995, Ann. of the Israel Physical Society, Vol. 11: Asymmetrical Planetary Nebulae, eds. A. Harpaz and N. Soker (Haifa, Israel), p. 240.
- Chevalier, R. A., & Luo D. 1994, ApJ, 421, 225 (CL).
- Chu, Y.-H., Chang, T. H., & Conway, G. M. 1997, ApJ, 482, 891.
- Dgani, R., Evans, N.J., & White, S.M. 1998, in preparation.
- Dyson J.E., 1968, Ap&SS, 1, 388.
- Garcia-Segura, G. 1997, ApJ, 489, L189.
- Jacoby, G. H. 1979, PASP, 91, 754.
- Jones, T. W., Kang, H., & Tregillis, I. L. 1994, ApJ, 432, 194.
- Jones, T. W., Ryu, D., & Tregillis, I. L. 1996, ApJ, 473, 365.
- Jun, B.I., & Jones, T. W. 1998, preprint.
- Jun, B.I., Jones, T. W., & Norman, M.L. 1996, ApJ, 468, L59.
- Kahn, F. D. 1969, Physica, 41, 172.
- Leuenhagen, U., Koesterke, L., & Hamann, W.-R., 1993, Acta Astron., 43, 329.
- Pacholczyk, A.G., 1970, Radio Astrophysics, San Francisco:Freeman
- Patriarchi, P., & Perinotto, M. 1991, A&AS, 91, 325.
- Soker, N. 1998, preprint.
- Zijlstra, A.A., Pottasch, S.R., & Bignell, C. 1989, A&AS, 79, 329.